

**Baxter**

# CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections

## in CLARITY Dual Chamber Container

### Description

CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections are sterile, nonpyrogenic, hypertonic solutions in a CLARITY Dual Chamber Container.

The sulfite-free Amino Acid Injections in the outlet port chamber are solutions of essential and nonessential amino acids.

The Dextrose Injections, USP in the injection port chamber are solutions for fluid replenishment and caloric supply.

After opening the seal between the chambers and mixing thoroughly, the admixed product is intended for intravenous use. See Table 1 for composition, pH, osmolarity, ionic concentration and caloric content of the admixed product.

The CLARITY Dual Chamber Container is a lipid-compatible plastic container (PL 2401 Plastic). The amount of water that can permeate from inside the container into the overwrap is insufficient to affect the solution significantly. Solutions in contact with the plastic container may leach out certain chemical components from the plastic in very small amounts; however, biological testing was supportive of the safety of the plastic container materials.

### Clinical Pharmacology

CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections administered intravenously provide biologically utilizable source material for protein synthesis and have value as a source of calories and water.

### Indications and Usage

CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections are indicated as a caloric component in a parenteral nutrition regimen and as the protein (nitrogen) source for offsetting nitrogen loss or for treatment of negative nitrogen balance in patients where: (1) the alimentary tract cannot or should not be used, (2) gastrointestinal absorption of protein is impaired, or (3) metabolic requirements for protein are substantially increased, as with extensive burns.

**Central Vein Administration:** Central vein infusion should be used when amino acid solutions are admixed with hypertonic dextrose to promote protein synthesis such as for hypercatabolic or depleted patients or those requiring long term parenteral nutrition.

**Peripheral Vein Administration:** For patients in whom the central vein route is not indicated, amino acid solutions diluted with low dextrose concentrations may be infused by peripheral vein.

### Contraindications

CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections are contraindicated in patients having intracranial or intraspinal hemorrhage, in patients who are severely dehydrated, in patients hypersensitive to one or more amino acids, and in patients with severe liver disease or hepatic coma.

Solutions containing corn-derived dextrose may be contraindicated in patients with known allergy to corn or corn products.

### Warnings

Additives may be incompatible including fat emulsions. Consult with pharmacist, if available. When introducing additives, use aseptic techniques. Mix thoroughly.

Because of the potential for life-threatening events, caution should be taken to ensure that precipitates have not formed in any parenteral nutrient admixture.

These CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections, **must be** admixed prior to infusion. For admixing instructions see **Directions for Use of Plastic Container**.

The infusion of hypertonic nutrient injections into a peripheral vein may result in vein irritation, vein damage, and thrombosis. After mixing, strongly hypertonic nutrient injections should only be administered through an indwelling intravenous catheter with the tip located in a large central vein, such as the superior vena cava.

Proper administration of these admixed amino acid/dextrose injections requires a knowledge of fluid and electrolyte balance and nutrition as well as clinical expertise in recognition and treatment of the complications which may occur.

### Laboratory Tests

**Frequent clinical evaluation and laboratory determinations are necessary for proper monitoring during administration.** Studies should include blood sugar, serum proteins, kidney and liver function tests, electrolytes, complete blood count with differential, carbon dioxide combining power or content, serum osmolarities, blood cultures, and blood ammonia levels.

Administration of amino acid solutions to a patient with hepatic insufficiency may result in serum amino acid imbalances, hyperammonemia, stupor, and coma.

**Pediatric Use:** Use of CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections in pediatric patients is governed by the same considerations that affect the use of any amino acid solution in pediatrics. The amount administered is dosed on the basis of grams of amino acids/kg of body weight/day. Two to 3 g/kg of body weight for infants with adequate calories are generally sufficient to satisfy protein needs and promote positive nitrogen balance. Solution administrations by peripheral vein should not exceed twice normal serum osmolarity (718 mOsmol/L).

**Central Vein Administration:** Hypertonic mixtures of amino acid/dextrose injections may be administered safely by continuous infusion through a central vein catheter with the tip located in the vena cava. In addition to meeting nitrogen needs, the administration rate is governed, especially during the first few days of therapy, by the patient's tolerance to dextrose, as indicated by frequent determinations of urine and blood sugar levels. Daily intake of amino acids in dextrose should be increased gradually to the maximum required dose.

Sudden cessation in administration of these admixed injections may result in insulin reaction due to continued endogenous insulin production. Parenteral nutrition mixtures should be withdrawn slowly.

**Peripheral Vein Administration:** For patients requiring parenteral nutrition in whom the central vein route is not indicated, low concentration amino acid/dextrose injections may be administered by peripheral vein. In pediatric patients, the final solution should not exceed twice normal serum osmolarity (718 mOsmol/L).

### Directions for Use of Plastic Container

**WARNING: Do not use plastic containers in series connections. Such use could result in air embolism due to residual air being drawn from the primary container before administration of the fluid from the secondary container is completed.**

**BE SURE THE CONTENTS OF BOTH CHAMBERS ARE MIXED TOGETHER AFTER OPENING SEAL BETWEEN CHAMBERS. After opening seal between chambers, lipids and/or additives can be introduced to the container. Thorough mixing ensures complete delivery of all ingredients.**

### To Open

Tear overwrap across top at slit and remove solution container. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually.

Check to ensure seal between chambers is intact, *i.e.*, solutions are contained in separate chambers. Check for minute leaks by separately squeezing each chamber. If external leaks or leakage between the chambers are found, discard solution as sterility or stability may be impaired.

### To Mix Solutions

Grasp the container firmly on each side of the top of the bag and roll bag to open seal between chambers as shown in Figure 1. Mix solutions thoroughly as shown in Figure 2. Check for leaks.

**Storage:** If removed from the overwrap and the contents are not mixed, CLINIMIX Injection solutions may be stored under refrigeration for up to 9 days.

Upon mixing of bag contents, CLINIMIX Injection solutions remain stable when stored under refrigeration, not to exceed 9 days from when the product was originally removed from the overwrap.




Figure 1



Figure 2

CLINIMIX Injection solutions containing additives should be used promptly after admixture. Any storage should be under refrigeration and limited to a brief period of time, less than 24 hours.

### To add Fat Emulsion for 3-in-1 admixture:

See **Warnings** section regarding incompatible additives including fat emulsions.

1. Prior to adding fat emulsion, mix amino acid and dextrose injection as shown in Figure 2.
2. Prepare fat emulsion transfer set following instructions provided.
3. Attach transfer set to fat emulsion bottle, using aseptic technique.
4. Twist off protector on the additive port of the CLARITY container.
5. Attach the transfer set to the exposed additive port.
6. Open clamp on transfer set.
7. After completing transfer, use appropriate plastic clamp or metal ferrule to seal off additive port tube.
8. Remove transfer set.
9. Mix contents of CLARITY container thoroughly. Check for leaks.

**Storage:** Storage of the 3-in-1 admixture must be under refrigeration and limited to a brief period of time, no longer than 24 hours. See **Warnings** section regarding incompatible additives.

### To Add Medication

**WARNING:** Additives may be incompatible.

Supplemental medication may be added with a 19 to 22 gauge needle through the medication port.

1. Prepare medication port.
2. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
3. Mix solution and medication thoroughly. For high density medication, such as potassium chloride, squeeze ports while ports are upright and mix thoroughly.
4. Check for leaks.

### Preparation for Administration

1. Suspend container from eyelet support.
2. Twist off protector from outlet port at bottom of container.
3. Attach administration set. Refer to complete directions accompanying set.

### How Supplied

See Table 1.

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. Protect from freezing. It is recommended that the product be stored at room temperature (25°C/77°F): brief exposure up to 40°C/104°F does not adversely affect the product.

Refrigerated storage is limited to 9 days once overwrap has been opened.

Do not use if overwrap has been previously opened or damaged.

### Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Printed in USA

Baxter, Clinimix, and Clarity are trademarks of Baxter International Inc.

07-19-57-358

Rev. July 2010

\*BAR CODE POSITION ONLY

071957358

**Metabolic:** The following metabolic complications have been reported: metabolic acidosis, hypophosphatemia, alkalosis, hyperglycemia and glycosuria, osmotic diuresis and dehydration, rebound hypoglycemia, elevated liver enzymes, hypo- and hypervitaminosis, electrolyte imbalances, and hyperammonemia. Frequent clinical evaluation and laboratory determinations are necessary, especially during the first few days of therapy to prevent or minimize these complications.

Caution must be exercised in the administration of these admixed amino acid/dextrose injections to patients receiving corticosteroids or corticotropin.

These admixed injections should be used with caution in patients with overt or known subclinical diabetes mellitus.

Drug product contains no more than 25 mcg/L of aluminum.

**Carcinogenesis, Mutagenesis, Impairment of Fertility:** Studies with CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections have not been performed to evaluate carcinogenic potential, mutagenic potential, or effects on fertility.

**Pregnancy:** Teratogenic Effects

Pregnancy Category C. Animal reproduction studies have not been conducted with CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections. It is also not known whether CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections should be given to a pregnant woman only if clearly needed.

**Nursing Mothers:** Caution should be exercised when CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections are administered to a nursing woman.

**Pediatric Use:** Dextrose is safe and effective for the stated indications in pediatric patients (see **Indications and Usage**). As reported in the literature, the dosage selection and constant infusion rate of intravenous dextrose must be selected with caution in pediatric patients, particularly neonates and low birth weight infants, because of the increased risk of hyperglycemia/hypoglycemia. Frequent monitoring of serum glucose concentrations is required when dextrose is prescribed to pediatric patients, particularly neonates and low birth weight infants.

Safety and effectiveness of CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections in pediatric patients have not been established by adequate and well-controlled studies. However, the use of amino acid injections in pediatric patients as an adjunct in the offsetting of nitrogen loss or in the treatment of negative nitrogen balance is referenced in the medical literature. See **Dosage and Administration**.

**Geriatric Use:** Clinical studies of CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from other younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or drug therapy.

### Adverse Reactions

See **Warnings and Precautions**

Too rapid infusion of these CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections may result in diuresis, hyperglycemia, glycosuria, and hyperosmolar coma. Continual clinical monitoring of the patient is necessary in order to identify and initiate measures for these clinical conditions.

Reactions that may occur because of the solution or the technique of administration include febrile response, infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection, extravasation, and hypervolemia. Policies and procedures should be established for the recognition and management of such reactions.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures, and save the remainder of the fluid for examination if deemed necessary.

### Dosage and Administration

If a patient is unable to take oral nourishment for a prolonged period of time, institution of total parenteral nutrition should be considered.

The total daily dose of CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections depends on the patient's metabolic requirement and clinical response. The determination of nitrogen balance and accurate daily body weights, corrected for fluid balance, are probably the best means of assessing individual nitrogen requirements.

Recommended Dietary Allowances\* of protein range from approximately 0.75 g/kg of body weight for adults to 1.68 g/kg for infants up to three months of age. It must be recognized, however, that protein as well as caloric requirements in traumatized or malnourished patients may be increased substantially. Daily amino acid doses of approximately 1.0 to 1.5 g/kg of body weight for adults with adequate calories are generally sufficient to satisfy protein needs and promote positive nitrogen balance.

For the initial treatment of trauma or protein calorie malnutrition, higher doses of protein with corresponding quantities of carbohydrate will be necessary to promote adequate patient response to therapy. The severity of the illness being treated is the primary consideration in determining proper dose level. Such higher doses, especially in infants, must be accompanied by more frequent laboratory evaluation.

Care should be exercised to insure the maintenance of proper levels of serum potassium. Quantities of 60 to 180 mEq of potassium per day have been used with adequate clinical effect. It may be necessary to add quantities of this electrolyte to these admixed injections, depending primarily on the amount of carbohydrate administered to and metabolized by the patient.

Patients receiving CLINIMIX sulfite-free (Amino Acid in Dextrose) Injections without electrolytes should be monitored frequently and their electrolyte requirements individualized.

Total daily fluid requirements can be met beyond the volume of amino acids solution by supplementing with noncarbohydrate or carbohydrate-containing electrolyte solutions.

Maintenance vitamins, additional electrolytes, and trace elements should be administered as required.

Table 1

How Supplied			Composition																	Caloric Content (kcal/L)							
After mixing, the product represents	1000 mL Code and NDC Number	2000 mL Code and NDC Number	Dextrose Hydrous, USP <sup>1</sup> (g/100 mL)	Amino Acids (g/100 mL)	Total Nitrogen (mg/100 mL)	Essential Amino Acids (mg/100 mL)										Nonessential Amino Acids (mg/100 mL)					Anion Profile (mEq/L) <sup>2</sup>		pH <sup>5</sup> (range)	Osmolarity (mOsmol/L) (calc)	From Dextrose	From Amino Acids	TOTAL (Dextrose and Amino Acids)
						Leucine - (CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> CH (NH <sub>2</sub> ) COOH	Isoleucine - CH <sub>3</sub> CH <sub>2</sub> CH (CH <sub>3</sub> ) CH (NH <sub>2</sub> ) COOH	Valine - (CH <sub>3</sub> ) <sub>2</sub> CHCH (NH <sub>2</sub> ) COOH	Lysine (added as the hydrochloride salt) - H <sub>2</sub> N (CH <sub>2</sub> ) <sub>4</sub> CH (NH <sub>2</sub> ) COOH	Phenylalanine - (C <sub>6</sub> H <sub>5</sub> ) CH <sub>2</sub> CH (NH <sub>2</sub> ) COOH	Histidine - (C <sub>3</sub> H <sub>3</sub> N <sub>2</sub> ) CH <sub>2</sub> CH (NH <sub>2</sub> ) COOH	Threonine - CH <sub>3</sub> CH (OH) CH (NH <sub>2</sub> ) COOH	Methionine - CH <sub>3</sub> S (CH <sub>2</sub> ) <sub>2</sub> CH (NH <sub>2</sub> ) COOH	Tryptophan - (C <sub>8</sub> H <sub>6</sub> N) CH <sub>2</sub> CH (NH <sub>2</sub> ) COOH	Alanine - CH <sub>3</sub> CH (NH <sub>2</sub> ) COOH	Arginine - H <sub>2</sub> NC (NH) NH (CH <sub>2</sub> ) <sub>3</sub> CH (NH <sub>2</sub> ) COOH	Glycine - H <sub>2</sub> NCH <sub>2</sub> COOH	Proline - [(CH <sub>2</sub> ) <sub>3</sub> NH CH] COOH	Serine - HOCH <sub>2</sub> CH (NH <sub>2</sub> ) COOH	Tyrosine - [C <sub>6</sub> H <sub>4</sub> (OH)] CH <sub>2</sub> CH (NH <sub>2</sub> ) COOH	Acetate <sup>3</sup>	Chloride <sup>4</sup>					
CLINIMIX 2.75/5 sulfite-free (2.75% Amino Acid in 5% Dextrose) Injection	Code 2B7725 NDC 0338-1132-03	Code 2B7701 NDC 0338-1083-04	5	2.75	454	201	165	160	159	154	132	116	110	50	570	316	283	187	138	11	24	11	6.0 (4.5 to 7.0)	525	170	110	280
CLINIMIX 4.25/5 sulfite-free (4.25% Amino Acid in 5% Dextrose) Injection	Code 2B7726 NDC 0338-1133-03	Code 2B7704 NDC 0338-1089-04	5	4.25	702	311	255	247	247	238	204	179	170	77	880	489	438	289	213	17	37	17	6.0 (4.5 to 7.0)	675	170	170	340
CLINIMIX 4.25/10 sulfite-free (4.25% Amino Acid in 10% Dextrose) Injection	Code 2B7727 NDC 0338-1134-03	Code 2B7705 NDC 0338-1091-04	10	4.25	702	311	255	247	247	238	204	179	170	77	880	489	438	289	213	17	37	17	6.0 (4.5 to 7.0)	930	340	170	510
CLINIMIX 4.25/20 sulfite-free (4.25% Amino Acid in 20% Dextrose) Injection	Code 2B7728 NDC 0338-1135-03	Code 2B7706 NDC 0338-1093-04	20	4.25	702	311	255	247	247	238	204	179	170	77	880	489	438	289	213	17	37	17	6.0 (4.5 to 7.0)	1435	680	170	850
CLINIMIX 4.25/25 sulfite-free (4.25% Amino Acid in 25% Dextrose) Injection	Code 2B7729 NDC 0338-1136-03	Code 2B7707 NDC 0338-1095-04	25	4.25	702	311	255	247	247	238	204	179	170	77	880	489	438	289	213	17	37	17	6.0 (4.5 to 7.0)	1685	850	170	1020
CLINIMIX 5/15 sulfite-free (5% Amino Acid in 15% Dextrose) Injection	Code 2B7730 NDC 0338-1137-03	Code 2B7709 NDC 0338-1099-04	15	5	826	365	300	290	290	280	240	210	200	90	1035	575	515	340	250	20	42	20	6.0 (4.5 to 7.0)	1255	510	200	710
CLINIMIX 5/20 sulfite-free (5% Amino Acid in 20% Dextrose) Injection	Code 2B7731 NDC 0338-1138-03	Code 2B7710 NDC 0338-1101-04	20	5	826	365	300	290	290	280	240	210	200	90	1035	575	515	340	250	20	42	20	6.0 (4.5 to 7.0)	1505	680	200	880
CLINIMIX 5/25 sulfite-free (5% Amino Acid in 25% Dextrose) Injection	Code 2B7732 NDC 0338-1139-03	Code 2B7711 NDC 0338-1103-04	25	5	826	365	300	290	290	280	240	210	200	90	1035	575	515	340	250	20	42	20	6.0 (4.5 to 7.0)	1760	850	200	1050

In many patients, provision of adequate calories in the form of hypertonic dextrose may require the administration of exogenous insulin to prevent hyperglycemia and glycosuria.

Fat emulsion administration should be considered when prolonged (more than 5 days) parenteral nutrition is required in order to prevent essential fatty acid deficiency (EFAD). Serum lipids should be monitored for evidence of EFAD in patients maintained on fat-free TPN.

Intravenous fat emulsions provide approximately 1.1 kcal per mL (10%), 2.0 kcal per mL (20%), or 3.0 kcal per mL (30%) and may be admixed along with amino acid/dextrose injections in the CLARITY Container to supplement caloric intake.

Depending upon the clinical condition of the patient, approximately 3 liters of solution may be administered per 24 hour period. When used postoperatively, the therapy should begin with 1000 mL on the first postoperative day. Thereafter, the dose may be increased to 3000 mL per day.

Do not administer unless seal between chambers is opened, other seals are intact, and solution is clear and thoroughly mixed.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit. Use of a final filter is recommended during administration of all parenteral solutions, where possible.


A slight yellow color does not alter the quality and efficacy of this product.

Additives may be incompatible. Complete information is not available. Those additives known to be incompatible should not be used. Consult with pharmacist, if available.

If, in the informed judgment of the physician, it is deemed advisable to introduce additives, use aseptic technique. Mix thoroughly when additives have been introduced.

Do not store solutions containing additives. These amino acid with electrolytes/dextrose with calcium injections should be used promptly after mixing. Any storage with additives should be under refrigeration and limited to a brief period of time, less than 24 hours.

\* Food and Nutrition Board National Academy of Sciences - National Research Council (Revised 1989).



- Balanced by ions from amino acids.
- Derived from glacial acetic acid (for pH adjustment).
- Contributed by the lysine hydrochloride.
- pH of sulfite-free Amino Acid Injection in the outlet port chamber was adjusted with glacial acetic acid.